

## Review Article

# Pharmacological Effects of Verticine: Current Status

Zhenhua Yin <sup>1,2</sup>, Juanjuan Zhang,<sup>1,2</sup> Qingfeng Guo,<sup>1,2</sup> Lin Chen,<sup>1,2</sup>  
Wei Zhang,<sup>1,2</sup> and Wenyi Kang <sup>1,2</sup>

<sup>1</sup>Zhengzhou Key Laboratory of Medicinal Resources Research, Huanghe Science and Technology College, Zhengzhou 450063, China

<sup>2</sup>Henan Joint International Research Laboratory of Drug Discovery of Small Molecules, Zhengzhou 450063, China

Correspondence should be addressed to Wenyi Kang; kangweny@hotmail.com

Received 21 November 2018; Revised 18 February 2019; Accepted 27 February 2019; Published 7 March 2019

Academic Editor: Nunziatina De Tommasi

Copyright © 2019 Zhenhua Yin et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Verticine is the major bioactive constituent of *Fritillaria* as a kind of Traditional Chinese Medicine. Pharmacological researches have reported various benefits of verticine, including anticancer, anti-inflammatory, protecting against acute lung injury, tracheobronchial relaxation, antitussive, expectorant, sedative, and analgesic activities, in addition to inhibiting proliferation of cultured orbital fibroblast, angiotensin converting enzyme (ACE), and acetylcholinesterase (AChE) and inhibiting hERG potassium channels. The underlying mechanisms of verticine are still under investigation. This review will comprehensively summarize the metabolism, biological activities, and possible mechanism of verticine.

## 1. Introduction

Verticine (Figure 1) belongs to a kind of isosterol alkaloid, is the major bioactive constituent of *Fritillaria* as Traditional Chinese Medicine that is widely used as an antitussive and expectorant [1]. Pharmacological researches on verticine have reported its valuable benefits in a variety of diseases, especially its anticancer effect. In this paper, the pharmacological effects, including metabolism, antitumor, anti-inflammatory, protection against acute lung injury diastolic bronchus, inhibition of angiotensin converting enzyme, and acetylcholinesterase, antitussive expectorant, sedative analgesia, were summarized, which provides theoretical references for its clinical application.

## 2. Metabolism

Pharmacokinetics of verticine is closely related to its biological activity, and the metabolism is influenced by the mode of administration, sex, and animal types. Pharmacokinetic behavior of verticine can provide the theory reference for clinical medicine.

In rabbits model, the pharmacokinetics of verticine was different between the intragastric (ig) administration and intravenous (iv) administration (Table 1). The  $t_{1/2}$  of ig

administration was three times longer than that of iv administration, suggesting that there might be a reabsorption process after ig administration. However, it showed a very low bioavailability of 10.65%, which might be its low solubility in water, incomplete absorption, or metabolism of gastrointestinal enzymes and efflux pumps [2].

The pharmacokinetics of verticine was influenced significantly by sex. Verticine was eliminated slowly in the plasma of male Sprague-Dawley rat but not in female rats, and gender-related differences were also observed significantly in the pharmacokinetic parameters (Table 1). Drug concentration in blood and tissue in male rats was significantly higher except for several tissues, such as fat, muscle, and skin (data not given). Urinary cumulative excretion of verticine in female rats ( $0.12 \pm 0.04\%$ ) was lower than that of male rats ( $0.90 \pm 0.28\%$ ), and fecal cumulative excretion between female rats ( $0.23 \pm 0.06\%$ ) and male rats ( $0.27 \pm 0.06\%$ ) had no difference. Differences of sex-associated metabolism for verticine in rats are mainly due to sex-dependent expression and activity of drug metabolism enzymes and P-glycoprotein (P-gp) [3]. In addition, the main pharmacokinetic parameters of verticine were obviously different from [3] in Sprague-Dawley rats' plasma after gastric gavage extract of *Fritillaria thunbergii* Miq. The V1/F was 40.832 L/mg, indicating that verticine was mainly distributed in blood, intracellular fluid,

TABLE 1: The pharmacokinetics of verticine in references.

Models	Extracts/compounds	Methods	Mode of administration	Pharmacokinetic parameters	Ref.
Rabbits	Verticine	LC-MS/MS	ig administration iv administration	A two-compartment model, $C_{max} = 48.31 \pm 74.0$ ng/mL, $AUC_{0-24} = 270.08 \pm 80.17$ ng/(mL/h), $t_{1/2} = 6.38 \pm 3.11$ h A three-compartment model, $C_{max} = 84.39 \pm 15.39$ ng/mL, $AUC_{0-24} = 50.72 \pm 14.02$ ng/(mL/h), $t_{1/2} = 2.19 \pm 1.07$ h	[2]
Sprague-Dawley female rats Sprague-Dawley male rats	4.25 g/kg <i>Fritillaria thunbergii</i> Miq. Extract	LC-MS/MS	a single oral administration	$C_{max} = 43.7 \pm 22.7$ ng/mL, $T_{max} = 1.5 \pm 0.7$ h, $t_{1/2} = 4.2 \pm 2.0$ h, $AUC_{0-24} = 214.2 \pm 84.6$ ng/(mL/h), $AUC_{0-24} = 214.3 \pm 84.5$ ng/(mL/h), $CL/F = 128.9 \pm 32.6$ L/kg/h, $V/F = 781.3 \pm 305.6$ L/kg $C_{max} = 57.6 \pm 21.6$ ng/mL, $T_{max} = 2.9 \pm 1.7$ h, $t_{1/2} = 6.2 \pm 1.9$ h, $AUC_{0-24} = 662.4 \pm 277.9$ ng/(mL/h), $AUC_{0-24} = 665.3 \pm 213.3$ ng/(mL/h), $CL/F = 41.5 \pm 20.1$ L/h/kg, $V/F = 374.1 \pm 186.2$ L/kg	[3]
Sprague-Dawley Rats	<i>Fritillaria thunbergii</i> Miq. Extract (0.45 g/kg body weight)	RRLC-MS/MS	gastric gavage	A two-compartment model, $C_{max} = 3.671 \pm 0.876$ $\mu$ g/L, $T_{max} = 32.5 \pm 11.292$ min, $t_{1/2\alpha} = 29.269 \pm 24.156$ min, $t_{1/2\beta} = 162.897 \pm 30.669$ min, $AUC_{0-24} = 628.568 \pm 100.99$ $\mu$ g/(L*min), $AUC_{0-24} = 630.875 \pm 102.136$ $\mu$ g/(L*min), $CL/F = 0.423 \pm 0.075$ L/min/kg, $V1/F = 40.832 \pm 17.616$ L/kg	[4]

\*  $p < 0.05$ , significantly different from female rats.




FIGURE 1: Chemical structure of verticine.

and extracellular fluid, and it was widely distributed *in vivo* [4].

Further study showed that the intestinal absorption of verticine involved both active transport, facilitated diffusion, and resulted in the low bioavailability in male and female rats [34]. Caco-2 cell monolayer exerted an effect on the intestinal absorption of verticine. Verticine transport was of concentration-dependent type, and both  $P_{app(AP-BL)}$  and  $P_{app(BL-AP)}$  were higher at 4°C than that at 37°C. When the P-glycoprotein (P-gp) inhibitors, verapamil, and cyclosporin A were present, the  $P_{app(AP-BL)}$  was higher and  $P_{app(BL-AP)}$  was lower, and the absorption permeability was not affected by EDTA- $Na_2$ . The P-gp inhibitors could increase the absorption of verticine, and EDTA- $Na_2$  had no discernible effect on absorption. The intestinal absorption of verticine across Caco-2 cell monolayers involved active transport rather than passive diffusion, and verticine was a substrate of P-gp [35].

### 3. Pharmacological Effects

**3.1. Antitumor Effect.** The treatment of cancer is mainly based on chemotherapy. Multiple drug resistance limited the

improvement of chemotherapy efficacy and also became an important reason for the recurrence and metastasis of cancer. Verticine had the effects of anti-cell proliferation and apoptosis in many human tumor cell lines and could reverse the multidrug resistance of some drug-resistant cell lines, such as breast cancer, leukemia, lung cancer, and gastric cancer cells.

**3.1.1. Anti-Breast Cancer Effect.** Verticine could inhibit the proliferation of breast cancer cell and induce its apoptosis and significant multidrug resistance reversal activity against breast cancer cell. Tong et al. confirmed that verticine had effects on inhibiting proliferation and reversing multidrug resistance against MCF-7/A [4]; the result was consistent with the research of Hu KW and Chen XY (1998) which confirmed that verticine showed significant multidrug resistance reversal activity against MCF-7/ADR and MCF-7 cell lines [5]. In recent years, researchers have focused on apoptosis-related protein Bcl-2. They found that Bcl-2 was involved in the tumor pathogenesis by inhibiting apoptosis. At the same time, some studies believed that the expression of Bcl-2 was one of the mechanisms of tumor resistance, and expression of Bcl-2 in tumor cells was detected during the treatment of triphenylamine (TAM) [36–38]. For breast cancer cell, Chen and Chen had proved that verticine could inhibit the proliferation of MCF-7/TAM cells ( $IC_{50} = 191.16$  g/mL at 48 h and  $IC_{50} = 138.10$  g/mL at 72 h) and induce its apoptosis by decreasing expression of Bcl-2 [6]. In addition, verticine showed distinct inhibiting effect on breast cancer 4T1 cells ( $IC_{50} = 14.7$   $\mu$ mol/L at 48 h); the mechanism might be related to downregulation of TGF- $\beta$ , VEGF, and MCP-1 secretion and decrease of TGF- $\beta$  and VEGF mRNA expression in 4T1 cells, thereby regulating its tumor inflammatory microenvironment to show the effect of anti-cancer [7]. Further, its

mechanism was investigated on 4T1 cells, and the results showed that verticine could regulate blood viscosity, improve blood flow state, reduce the expression of u-PA, VEGF, and PAI-1 protein and the secretion of IL-8, reduce the infiltration of neutrophils, improve TFPI-2 protein expression to promote tumor apoptosis, and inhibit angiogenesis and reduced cell transfer rate [8].

**3.1.2. Anti-Human Leukemia Effect.** Verticine could inhibit the proliferation of human leukemia cell and induce apoptosis of multidrug resistant leukemia; the mechanism was likely to be related to protein expression, redox imbalance, and caspase-3. In the early studies, verticine could inhibit the proliferation of HL-60 and K562 and reverse the multidrug resistance reversal activity against HL-60/ADR and K562/A02, which might be the increase of intracellular drug concentration and inhibition of P-gp protein expression in drug-resistant cells [5, 9]. ROS was an important signal molecule in cells, involved in many events, for example, cell proliferation, apoptosis, and multidrug resistance [39, 40]. It induced ROS explosion and reduced GSH content in tumor cell to inhibit tumor cell proliferation and induce apoptosis, which had an antitumor effect [41, 42]. This finding was consistent with that of Qi et al. (2017) who proved the effect of verticine on cell viability, proliferation, and apoptosis of human leukemia and the function of reactive oxygen species and redox imbalance in this progress [10, 11]. Some alkaloids could activate caspase-3 and caspase-dependent cell apoptosis [43–45], and stimulating the ROS production of K562 cell could promote caspase-3 expression and induce apoptosis [46, 47]. Therefore, verticine might also activate caspase-3-related pathway in the process of stimulating ROS-induced apoptosis in K562/A02 cells.

**3.1.3. Anti-Lung Cancer Effect.** Lung resistance protein (LRP) was closely related to primary resistance to cisplatin (DDP). Excision repair cross-complement 1 (ERCC1) mRNA enhanced DNA repair capacity to mediate multidrug resistance of platinum drugs [48]. Tang et al. found that verticine could inhibit A549/DDP cell proliferation in a dose-dependent manner within 48 h treatment, and the reverse index was 3.73. After 48 h, cell apoptosis rate and LRP positive cell of verticine were  $38.16 \pm 2.25$  and  $(5.8 \pm 1.3)/HP$ , respectively. In addition, verticine could obviously decreased the expression levels of ERCC1 mRNA and LRP. It indicated that verticine could reverse MDR of A549/DDP cell line. Its mechanism might be apoptosis induction and downregulated expression of LRP and ERCC1 mRNA [12, 13].

**3.1.4. Anti-Gastric Cancer Effect.** Verticine could inhibit proliferation of SGC-7901 and SGC-7901/VCR, but had no obvious multiple drug resistance reversal effect [14].

**3.2. Anti-Inflammatory Effect.** Inflammation is a complex biological response mediated by activated inflammatory cells and immunocytes, involving a balance between proinflammatory and anti-inflammatory factors [15, 49]. Verticine showed anti-inflammatory effect. Zhang et al. proved that

verticine could regulate inflammatory microenvironment of 4T1 breast cancer cell by controlling the release of inflammatory factors and decreasing the expression of mRNA [7]. Additionally, verticine could inhibit the gene and protein expression of MUC5AC mucin induced by EGF, PMA, or TNF- $\alpha$ , by directly acting on airway epithelial cells, and the production of MUC5AC mucin protein induced by EGF, PMA, or TNF- $\alpha$ . This finding was consistent with the traditional use of *F. thunbergii* as remedy for diverse inflammatory pulmonary diseases [16]. At the same time, Yi et al. confirmed that verticine significantly inhibited tumor necrosis factor (TNF)- $\alpha$ , interleukin (IL)-6, and IL-1 $\beta$ , increased IL-10 production in lipopolysaccharide (LPS)-stimulated RAW 264.7 macrophages, and inhibited the phosphorylation of p38, ERK and c-Jun N-terminal kinase (JNK) as well as decreased p65 and I $\kappa$ B, which indicated that verticine inhibited the production of inflammatory cytokines induced by LPS through blocking MAPKs and NF- $\kappa$ B signaling pathways [50]. Verticine could inhibit the production of proinflammatory cytokines, such as IL-6, IL-8, and TNF- $\alpha$ , reducing MAPKs phosphorylation and the nuclear NF- $\kappa$ B expression in PMACI-induced HMC-1 [17].

In addition, the activity of T cell is inhibited, which can also achieve anti-inflammatory effect; KV1.3 potassium channels play a key role in the activation of T cells. Verticine could inhibit Kv1.3 channels in a concentration-dependent manner ( $IC_{50}=142.1 \mu M$  at 150 ms) [18].

**3.3. Protection against Acute Lung Injury.** The pathogenesis of acute lung injury was complex, but its essence was the damage of lung endothelial cells and alveolar epithelial cells caused by excessive inflammation [51, 52]. LPS can activate and amplify inflammatory reactions in the body, causing the accumulation of inflammatory cells in the lungs [53]. Verticine had protective effect on LPS-induced ALI in mice; the mechanism was related to the inhibition of the inflammatory factors, the downregulation of the phosphorylation level of MAPKs in the inflammatory response signaling pathway, and the reduction of NF- $\kappa$ B gene transcriptional intensity [19–23].

**3.4. Tracheobronchial Relaxation and Antitussive Effects.** Tracheal bronchial relaxation of verticine could be attributed to M receptor and calcium ions. Verticine showed strong inhibitory effect on the contraction of isolated tracheal strips of guinea pigs induced by carbachol; the results implied that the effect could be attributed to M receptor of the tracheal wall [24]. This result was consistent with that of Chan et al. who demonstrated the mechanisms of competitive antagonism of muscarinic pathway and also the inhibition of influx of calcium ions [25]. At the same time, verticine could also significantly elevate the concentration of cAMP in the HEK cells transfected with muscarinic M<sub>2</sub> receptor plasmid [54]. However, verticine did not exhibit agonistic  $\beta_2$  receptor activity. It could be seen that the effect of verticine on diastolic bronchus was not produced by the agonist  $\beta_2$  receptor [55].

Generally, M receptor is inhibited, which could have a certain antitussive effect, and verticine is an active constituent

TABLE 2: The biological activities of verticine in references.

Activities	Models	Biological activities	Action mechanism	Ref.
Anti-breast cancer	MCF-7	Inhibit proliferation, reversing multidrug resistance	-	[4, 5]
	MCF-7/TAM	Inhibit proliferation (at 48 h $IC_{50}=191.16$ g/mL; at 72 h, $IC_{50}=138.10$ g/mL), induce apoptosis	Decrease expression of Bcl-2	[6]
	4T1	Inhibit proliferation (at 48h, $IC_{50}=14.7$ $\mu$ mol/L)	(1) Down-regulate TGF- $\beta$ , VEGF and MCP-1 secretion, decrease TGF- $\beta$ and VEGF mRNA expression, regulating its tumor inflammatory microenvironment. (2) Regulate blood viscosity, improve blood flow state, reduce the expression of u-PA, VEGF, PAI-1 protein and the secretion of IL-8, reduce the infiltration of neutrophils, improve TFPI-2 protein expression	[7, 8]
Anti-human leukemia	HL60, HL-60/ADR, K562 K562/A02	Inhibit proliferation ( $IC_{50}=288.27\pm34.23$ , $256.52\pm26.15$ , $320.80\pm36.52$ , $300.06\pm33.18$ , $\mu$ g/mL), reverse multidrug resistance	Increase intracellular drug concentration and inhibit P-gp protein expression	[5, 9]
	K562/A02	Inhibit the cell viability and induce apoptosis, different concentrations of verticine (100, 200, 400 mol/L); the cell viabilities were $0.392\pm0.040$ , $0.314\pm0.022$ , $0.161\pm0.033$	Induce the ROS outbreak and increase the GSH content, redox imbalance	[10, 11]
	K562	Inhibit proliferation ( $IC_{50}=238$ mol/L)		
Anti-lung cancer	A549/DDP	Inhibit proliferation, induce apoptosis, reversing multidrug resistance	Down-regulate expression of LRP and ERCC1 mRNA	[12, 13]
Anti-gastric cancer	SGC-7901 and SGC-7901/VCR	Inhibit proliferation	-	[14]
Anti-inflammatory effect	4T1	Regulate inflammatory microenvironment	Control release of inflammatory factors, such as TGF- $\beta$ , VEGF, MMP-9, and MCP-1, decreasing the expression of TGF- $\beta$ and VEGF mRNA	[6]
	Confluent NCI-H292 cells	Remedy for inflammatory pulmonary diseases	Inhibit gene and protein expression of MUC5AC mucin induced by EGF, PMA or TNF- $\alpha$ by directly acting on airway epithelial cells	[15]
	LPS-induced RAW264.7 macrophages	Inhibit production of inflammatory cytokines induced by LPS	Block MAPKs and NF-kB signaling pathways	[16]
	HMC-1 Cells HEK 293	Inhibit production of inflammatory cytokines anti-inflammatory	Regulate the Phosphorylation of NF- $\kappa$ B and MAPKs Inhibit Kv1.3 channels	[17] [18]
Protection against acute lung injury	Mice	Protective effect on acute lung injury	Inhibit expression of TNF- $\alpha$ , IL-2, IL-6 and IL-8 and COX-2, promote the synthesis and release of SP-A, decrease the levels of PGE2 and NO. And, in addition, down-regulate phosphorylation level of MAPKs in the inflammatory response signaling pathway, and reduce NF- $\kappa$ B gene transcriptional intensity	[19-23]

TABLE 2: Continued.

Activities	Models	Biological activities	Action mechanism	Ref.
Tracheobronchial Relaxation	Isolated tracheal strips of guinea pigs rat isolated tracheal and bronchial	Inhibit contraction Inhibit carbachol-induced contraction	M receptor Inhibit influx of calcium ions	[24] [25]
Antitussive effect	Mice	Antitussive effect for ammonium hydroxide induced cough (4 mg/kg) Antitussive effect for mechanical stimulation induced cough (4 mg/kg)	- -	[26] [26]
	Guinea pig trachea	Antitussive effect electrical stimulation induced cough (4 mg/kg)	-	[26]
	Cat superior laryngeal nerve	Inhibit cough frequency and increase latent period of cough induced by ammonia	-	[27]
Expectorant effect	Mice's tracheal	Enhance mice's tracheal phenol red output in expectorant evaluation	Increase tracheobronchial mucus secretion and decrease the viscosity of mucus	[27, 28]
Sedative effect	Mice	Reduce spontaneous activity, prolong pentobarbital sleep time and increase sleep rate	-	[29]
Analgesic effect	Mice	Inhibit writhing reaction induced by acetic acid	Block Nav1.7 ion channel ( $IC_{50}=47.2\pm 3.3 \mu M$ )	[17, 26]
Inhibit proliferation of cultured Orbital fibroblast	Thyroid-associated ophthalmopathy (TAO)-patients	Different concentrations (5, 25, 50, 75, 100 mg/L) inhibit the proliferation of orbital fibroblasts in vitro in a dose-dependent manner.	-	[30]
Inhibit angiotensin converting enzyme (ACE) activity	Rat plasma	Inhibit ACE I activity in a dose-dependent manner ( $IC_{50}=312.8 \mu M$ )	-	[31]
Inhibit acetylcholine (AChE) inhibitory activity	HEK293 cell line	At 100 $\mu g/mL$ , inhibition rate was only 25%	-	[32]
Inhibit hERG potassium channels	HEK293 cell line	Inhibit hERG peak tail currents with $IC_{50}$ value of 43.7 mM	channel inactivation, Mutation of Y652 to Alanine reduced sensitivity	[33]

of *Fritillaria* in relieving cough, which is related to acting on M receptor of tracheal smooth muscle to relax trachea and relieve tracheal spasm. For ammonium hydroxide induced cough in mice, mechanical stimulation of guinea pig trachea induced cough in guinea pigs, and electrical stimulation of cat superior laryngeal nerve induced cough in cats; verticine (4 mg/kg) had obvious antitussive effect. For mechanical stimulation induced cough in guinea pigs, the antitussive effect of verticine reached the peak at 30~60 min, and the antitussive effect could be sustained for about 1 h [26]. In addition, verticine could also significantly inhibit cough frequency and increase latent period of cough in mice induced by ammonia [27].

**3.5. Expectorant and Sedative Effects.** Verticine had expectorant and sedative effects. The effect was related to its ability to increase tracheobronchial mucus secretion and decrease the viscosity of mucus [27, 28]. Verticine could reduce the spontaneous activity of mice, inhibit the increasing of number of activities caused by caffeine, prolong the pentobarbital sleep time, and increase the sleep rate in mice [29].

**3.6. Analgesic Effect.** Verticine could inhibit writhing reaction induced by acetic acid in mice; at the concentration of 1 mg/kg, it had significant analgesic effect ( $P < 0.05$ ), and at the concentration of 2 mg/kg, it had very significant analgesic effect ( $P < 0.01$ ) [26]. Previous studies showed that traditional local anesthetics could play an analgesic role by nonselectively blocking the voltage-gated sodium channel subfamily, and selective Nav1.7 inhibitors were also demonstrated to be analgesic in animal models. Nav1.7 emerged as a potential target for the treatment of pain [56]. Verticine was able to block the Nav1.7 ion channel ( $IC_{50} = 47.2 \pm 3.3 \mu M$ ), which might be the analgesic mechanism of verticine [17].

**3.7. Other Effects.** Several studies showed that verticine also had other effects. Li et al. proved that verticine could inhibit the proliferation of cultured orbital fibroblast of the thyroid-associated ophthalmopathy (TAO)-patients; the inhibitory effect was obviously better than that of the normal [30]. Verticine could inhibit the activity of ACE I in a dose-dependent manner ( $IC_{50} = 312.8 \mu M$ ), which might be responsible, at least in part, for the antihypertensive action [31]. Unfortunately it showed no appreciable AChE inhibitory activity at a concentration of 100  $\mu g/mL$ , and the inhibition rate was only 25% [32]. In addition, verticine could inhibit the hERG peak tail currents with  $IC_{50}$  value of 43.7 mM, and multiple results suggested that the inhibition was related to the channel inactivation. Further investigation showed that the mechanism of the inhibition was related to the mutation of Y652 to Alanine reduced sensitivity to verticine, which suggested that Y652 was an important binding site of hERG for verticine [33]. The main concern for cardiac safety determination is the possible inhibition of hERG ion channels. So, verticine should be used with caution to avoid its toxic effect on the heart.

## 4. Conclusions

The summary of metabolism and pharmacological researches of verticine shows that the mode of administration, dose, and gender have effects on metabolism, and verticine possesses multiple pharmacological effects that are summarized in Table 2. Verticine can control the expression of related proteins, inhibit inflammatory factors, and destroy redox balance to achieve antitumor effect. The mechanism of anti-inflammatory effect was related to MAPKs and NF- $\kappa$ B signaling pathways, and the protection against acute lung injury has close relation with anti-inflammatory effect. Acting on M receptor and inhibiting influx of calcium ions could inhibit tracheobronchial contraction, so as to play a role in relieving cough. The clinical studies of verticine remain somewhat elusive and show the risks of heart-safety.

In summary, verticine is a new potential plant-origin drug that has antitumor, anti-inflammation, protecting liver injury, and antitussive effect. The clinical effect should be focused on.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

This work was supported by the National Natural Science Foundation of China (31501552), Henan Province Industry-University-Research Cooperation Project (182107000033), and Key Research Projects of Colleges and Universities in Henan province (18A360019 and 19B360007).

## References

- [1] M. H. Chen, H. C. Wang, Y. M. Zhu et al., "Pharmacological research on *Fritillaria*," *Anhui Agricultural Science Bulletin*, vol. 13, no. 1, pp. 103–105, 2007.
- [2] L. L. Liu, L. H. Chen, W. F. Zhu et al., "Pharmacokinetics of peimine in rabbits," *Chinese Journal of Pharmaceuticals*, vol. 42, no. 12, pp. 914–916, 2011.
- [3] L.-H. Chen, H.-M. Zhang, Z.-Y. Guan et al., "Sex dependent pharmacokinetics, tissue distribution and excretion of peimine and peiminine in rats assessed by liquid chromatography-tandem mass spectrometry," *Journal of Ethnopharmacology*, vol. 145, no. 1, pp. 77–84, 2013.
- [4] X. L. Tong, *Reversing Multi-drug Resistance on Tumour Cells and Pharmacokinetics Study on Ingredients in a Traditional Chinese medicine *Fritillaria thunbergii* Miq*, Yunnan University of Traditional Chinese Medicine, 2016.
- [5] K. W. Hu and X. Y. Chen, "A screening study on the effects of activity ingredients in hinese herbs on anti-multidrug resistance cell line in vitro," *China Journal of Traditional Chinese Medicine and Pharmacy*, vol. 13, no. 2, pp. 10–12, 1998.
- [6] H. Y. Chen and X. Y. Chen, "Effect of peimine on inhibiting the MCF-7/TAM proliferation of human breast cancer cell and its influence on cell apoptosis," *Acta Chinese Medicine and Pharmacology*, vol. 40, no. 4, pp. 12–15, 2012.
- [7] Y. R. Zhang, H. S. Lin, and Y. Zhang, "Intervention effect of peimine, peiminine on the inflammatory microenvironment of

- 4T1 breast cancer cell," *Chinese Journal of Basic Medicine in Traditional Chinese Medicine*, vol. 20, no. 11, pp. 1504–1506, 2014.
- [8] N. Li, "Effect of Fuzheng Huoxue Jiedu Decoction on Tumor Stem Cells-Dependent PMNs on Tumor Metastasis, China Academy of Chinese Medical Sciences, 2017.
- [9] K. W. H. X. Zheng, J. Qi et al., "Study on reversing hematological cells multidrug resistance by verticine," *American Journal of Hematology*, vol. 20, no. 12, pp. 33–34, 1999.
- [10] Y. Qi, B. Liao, C. B. Xu et al., "Effects of peimine on cell viability and apoptosis of multidrug resistant human leukemia cells," *Shandong Medical Journal*, vol. 57, no. 26, pp. 17–20, 2017.
- [11] Y. Qi, B. Liao, C. B. Xu et al., "Peimine inhibits cell proliferation in human leukemia cell line K562 through redox imbalance," *China Medical Herald*, vol. 14, no. 18, pp. 16–19, 2017.
- [12] X. Y. Tang and Y. X. Tang, "Efficiency and mechanisms of peimine reversing multi-drug resistance of A549/DDP cell line," *Shandong Medical Journal*, vol. 52, no. 18, pp. 4–6, 2012.
- [13] X. Y. Tang, Y. X. Tang, and P. Xu, "Effect of peimine on ERCC1 mRNA and LRP expressions of A549/DDP multidrug resistance cell line," *Chinese Journal of Integrated Traditional and Western Medicine*, vol. 35, no. 12, pp. 1490–1493, 2015.
- [14] Z. Y. Gu, P. Zhang, Y. Z. Nie et al., "Screening for five alkaloids as reversal agents against gastric cancer multiple drug resistance and study on their mechanism," *Chinese Traditional and Herbal Drugs*, vol. 43, no. 6, pp. 1151–1156, 2012.
- [15] J. J. Worthington, T. M. Fenton, B. I. Czajkowska, J. E. Klementowicz, and M. A. Travis, "Regulation of TGF $\beta$  in the immune system: an emerging role for integrins and dendritic cells," *Immunobiology*, vol. 217, no. 12, pp. 1259–1265, 2012.
- [16] E. J. Kim, Y. P. Yoon, K. W. Woo et al., "Verticine, ebeiedine and suchengbeisine isolated from the bulbs of *Fritillaria thunbergii* inhibited the gene expression and production of MUC5AC mucin from human airway epithelial cells," *Phytomedicine*, vol. 23, no. 2, pp. 95–104, 2016.
- [17] J. Park, B. Lee, H. Kim et al., "Peimine inhibits the production of proinflammatory cytokines through regulation of the phosphorylation of NF- $\kappa$ B and MAPKs in HMC-1 Cells," *Pharmacognosy Magazine*, vol. 13, no. Suppl 2, pp. S359–S364, 2017.
- [18] J. Xu, W. Zhao, L. Pan et al., "Peimine, a main active ingredient of *Fritillaria*, exhibits anti-inflammatory and pain suppression properties at the cellular level," *Fitoterapia*, vol. 111, pp. 1–6, 2016.
- [19] G. X. Gui, X. C. Xia, Y. H. Ma et al., "Effect of Peimine on the expression of inflammatory factors in mice with endotoxin-induced acute lung injury," *Chinese Journal of Gerontology*, vol. 35, no. 18, pp. 5089–5091, 2015.
- [20] G. X. Gui, "Protective effect of peimine on acute lung injury in mice," *Clinical Journal of Chinese Medicine*, vol. 8, no. 26, pp. 4–6, 2016.
- [21] G. X. Gui, "An effect of the Peimine on COX-2, PGE2 and NO expression of mice with acute lung injury," *Clinical Journal of Chinese Medicine*, vol. 8, no. 34, pp. 11–13, 2016.
- [22] G. X. Gui, "An effect of PM on expression of TNF- $\alpha$ , IL-10 and PGE2 in mice with acute lung injury," *Clinical Journal of Chinese Medicine*, vol. 9, no. 1, pp. 1–4, 2017.
- [23] G. X. Gui, "An effect of Peimine on TNF- $\alpha$  Signaling Pathway in mice with endotoxin-induced acutelung injury," *Chinese Manipulation and Rehabilitation Medicine*, no. 4, pp. 75–78, 2017.
- [24] Y. Zhou, H. Ji, P. Li, and Y. Jiang, "Antimuscarinic function of five fritillaria alkaloids on guinea pig tracheal strips," *Journal of China Pharmaceutical University*, vol. 34, no. 1, pp. 58–60, 2003.
- [25] S.-W. Chan, P. Li, Y.-W. Kwan, and G. Lin, "In vitro tracheo-bronchial relaxation of *Fritillaria* alkaloids," *Chinese Journal of Natural Medicines*, vol. 9, no. 5, pp. 345–353, 2011.
- [26] B. C. Qian and H. J. Xu, "Studies on the antitussive and sedative activities of peimine and peiminine," *Acta Pharmaceutica Scientia*, vol. 20, no. 4, pp. 306–308, 1985.
- [27] D. Wang, J. Zhu, S. Wang et al., "Antitussive, expectorant and anti-inflammatory alkaloids from *Bulbus Fritillariae Cirrhosae*," *Fitoterapia*, vol. 82, no. 8, pp. 1290–1294, 2011.
- [28] B. Q. Lin, P. B. Li, Y. G. Wang et al., "The expectorant activity of naringenin," *Pulmonary Pharmacology & Therapeutics*, vol. 21, no. 2, pp. 259–263, 2008.
- [29] G. Q. Zhao, B. Ren, X. P. Dong et al., "The research progress of Chuanbeimu," *Pharmacy and Clinics of Chinese Materia Medica*, vol. 3, no. 6, pp. 59–64, 2012.
- [30] B. Li, Y. L. Zheng, and W. Z. Liu, "Influence of the *Fritillaria thunbergii*'s extract on the proliferation of the cultured Orbital fibroblast of tTAO-patients," *International Journal of Ophthalmology*, vol. 10, no. 2, pp. 235–237, 2010.
- [31] H. Oh, D.-G. Kang, S. Lee, Y. Lee, and H.-S. Lee, "Angiotensin converting enzyme (ACE) inhibitory alkaloids from *Fritillaria ussuriensis*," *Planta Medica*, vol. 69, no. 6, pp. 564–565, 2003.
- [32] Z.-D. Yang and D.-Z. Duan, "A new alkaloid from *Fritillaria ussuriensis* Maxim," *Fitoterapia*, vol. 83, no. 1, pp. 137–141, 2012.
- [33] L. Kan, W. Zhao, L. Pan et al., "Peimine inhibits hERG potassium channels through the channel inactivation states," *Biomedicine & Pharmacotherapy*, vol. 89, pp. 838–844, 2017.
- [34] Z.-Y. Guan, L.-H. Zhang, L.-H. Chen, W.-F. Zhu, and H.-N. Liu, "Rat intestinal absorption trait of peimine and peiminine in Thunberg Fritillary Bulb extract," *Acta Pharmaceutica Sinica*, vol. 48, no. 12, pp. 1836–1843, 2013.
- [35] L. Chen, X. Lu, X. Liang et al., "Mechanistic studies of the transport of peimine in the Caco-2 cell model," *Acta Pharmaceutica Sinica B (APSB)*, vol. 6, no. 2, pp. 125–131, 2016.
- [36] L. J. Wang, L. M. Sun et al., "Prognostic significance of the expression of apoptosis and bcl-2 protein in lung cancer," *Chinese Journal of Cancer Prevention and Treatment*, vol. 13, no. 4, pp. 290–293, 2006.
- [37] N. Takahashi, M. Yanagihara, Y. Ogawa, B. Yamanoha, and T. Andoh, "Down-regulation of Bcl-2-interacting protein BAG-1 confers resistance to anti-cancer drugs," *Biochemical and Biophysical Research Communications*, vol. 301, no. 3, pp. 798–803, 2003.
- [38] D. M. Cittelly, P. M. Das, V. A. Salvo, J. P. Fonseca, M. E. Burow, and F. E. Jones, "Oncogenic HER2 $\Delta$ 16 suppresses miR-15a/16 and deregulates BCL-2 to promote endocrine resistance of breast tumors," *Carcinogenesis*, vol. 31, no. 12, pp. 2049–2057, 2010.
- [39] E. Kalinina, M. Novichkova, N. P. Scherbak, V. Solomka, and A. N. Saprin, "GSH-dependent redox regulation and antioxidant enzymes in the formation of resistance to doxorubicin in K562 human erythroleukemia cells," *Advances in Experimental Medicine and Biology*, vol. 500, pp. 241–244, 2001.
- [40] J. Yao, X. Wei, and Y. Lu, "Chaetominine reduces MRP1-mediated drug resistance via inhibiting PI3K/Akt/Nrf2 signaling pathway in K562/Adr human leukemia cells," *Biochemical and Biophysical Research Communications*, vol. 473, no. 4, pp. 867–873, 2016.
- [41] L. Khoshtabiat, M. Mahdavi, G. Dehghan et al., "Oxidative stress-induced apoptosis in chronic myelogenous leukemia K562 cells by an active compound from the dithio-carba mate

- family,” *Asian Pacific Journal of Cancer Prevention*, vol. 17, no. 9, pp. 4267–4273, 2016.
- [42] H. Akçakaya, S. Tok, F. Dal, S. A. Cinar, and R. Nurten, “ $\beta$ -carotene treatment alters the cellular death process in oxidative stress-induced K562 cells,” *Cell Biology International*, vol. 41, no. 3, pp. 309–319, 2017.
- [43] D. Wang, Y. Jiang, K. Wu, S. Wang, and Y. Wang, “Evaluation of antitumor property of extracts and steroidal alkaloids from the cultivated *Bulbus Fritillariae ussuriensis* and preliminary investigation of its mechanism of action,” *BMC Complementary and Alternative Medicine*, vol. 15, no. 1, p. 29, 2015.
- [44] D.-D. Wang, Y. Feng, Z. Li et al., “In vitro and in vivo antitumor activity of *Bulbus Fritillariae cirrhosae* and preliminary investigation of its mechanism,” *Nutrition and Cancer*, vol. 66, no. 3, pp. 441–452, 2014.
- [45] D. Wang, S. Wang, Y. Feng et al., “Antitumor effects of *Bulbus Fritillariae cirrhosae* on Lewis lung carcinoma cells in vitro and in vivo,” *Industrial Crops and Products*, vol. 54, pp. 92–101, 2014.
- [46] X.-D. Wang, C.-Y. Li, M.-M. Jiang et al., “Induction of apoptosis in human leukemia cells through an intrinsic pathway by cathachunine, a unique alkaloid isolated from *Catharanthus roseus*,” *Phytomedicine*, vol. 23, no. 6, pp. 641–653, 2016.
- [47] J. Y. Luo, W. J. Kong, and M. H. Yang, “HJC, A new arylnaphthalene lignan isolated from *Justicia procumbens*, causes apoptosis and caspase activation in K562 leukemia cells,” *Journal of Pharmacological Sciences*, vol. 125, no. 4, pp. 355–363, 2014.
- [48] C. L. Tang, H. P. Yang, and X. D. Zhou, “Advances of DNA damage repair and cisplatin resistance mechanisms in lung cancer,” *Chinese Journal of Lung Cancer*, vol. 14, no. 12, pp. 960–964, 2011.
- [49] A. Minagar, P. Shapshak, R. Fujimura, R. Ownby, M. Heyes, and C. Eisdorfer, “The role of macrophage/microglia and astrocytes in the pathogenesis of three neurologic disorders: HIV-associated dementia, Alzheimer disease, and multiple sclerosis,” *Journal of the Neurological Sciences*, vol. 202, no. 1–2, pp. 13–23, 2002.
- [50] P. Yi, Y. Wu, H. Dong et al., “Peimine impairs pro-inflammatory cytokine secretion through the inhibition of the activation of NF- $\kappa$ B and MAPK in LPS-induced RAW264.7 macrophages,” *Immunopharmacology and Immunotoxicology*, vol. 35, no. 5, pp. 567–572, 2013.
- [51] P. Valta, A. Uusaro, S. Nunes, E. Ruokonen, and J. Takala, “Acute respiratory distress syndrome: frequency, clinical course, and costs of care,” *Critical Care Medicine*, vol. 27, no. 11, pp. 2367–2374, 1999.
- [52] U. J. Sachs, “Recent insights into the mechanism of transfusion-related acute lung injury,” *Current Opinion in Hematology*, vol. 18, no. 6, pp. 436–442, 2011.
- [53] Y. Fang, P. Xu, C. Gu et al., “Ulinastatin improves pulmonary function in severe burn-induced acute lung injury by attenuating inflammatory response,” *Journal of Trauma - Injury Infection and Critical Care*, vol. 71, no. 5, pp. 1297–1304, 2011.
- [54] Y. Zhou, H. Ji, B.-Q. Lin, Y. Jiang, and P. Li, “The effects of five alkaloids from *Bulbus Fritillariae* on the concentration of cAMP in HEK cells transfected with muscarinic M2 receptor plasmid,” *American Journal of Chinese Medicine*, vol. 34, no. 5, pp. 901–910, 2006.
- [55] X. Y. Yan, B. F. Cheng, H. H. Hou et al., “Evaluation and screening of four heterologous alkaloids stimulating  $\beta$ 2 receptor activity,” *Lishizhen Medicine and Materia Medica Research*, vol. 23, no. 7, pp. 1630–1631, 2012.
- [56] X. N. Liang, G. Yu, Z. B. Zheng et al., “Relationships between sodium channel NaV 1.7 and pain,” *Chinese Journal of Pharmacology and Toxicology*, vol. 29, no. 2, pp. 297–301, 2015.